

## Rab3

Cat.No. 107 011; Monoclonal mouse antibody, 100 µg purified IgG (lyophilized)

### Data Sheet

|                            |   |
|----------------------------|---|
| Reconstitution/<br>Storage | 100 µg purified IgG, lyophilized. For <b>reconstitution</b> add 100 µl H <sub>2</sub> O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet. |
| Applications               | <b>WB:</b> 1 : 1000 (AP staining)<br><b>IP:</b> not recommended<br><b>ICC:</b> 1 : 100 up to 1 : 500<br><b>IHC:</b> yes<br><b>IHC-P:</b> 1 : 200 up to 1 : 500<br><b>EM:</b> yes  |
| Clone                      | 42.1  |
| Subtype                    | IgG1 (κ light chain)  |
| Immunogen                  | Full length rat recombinant Rab3a (UniProt Id: P63012)  |
| Epitop                     | AA 95 to 151 from rat Rab3a (UniProt Id: P63012)  |
| Reactivity                 | Reacts with: human (P20336, P20337, Q96E17, O95716), rat (P63012, Q63941, P62824, Q63942), mouse (P63011, Q9CZT8, P62823, P35276), vertebrates, zebrafish (Q6DHJ2).<br>Other species not tested yet.  |
| Specificity                | Specific for all four Rab 3 isoforms a, b, c d.   |

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

## Background

Rab3 proteins belong to the Rab family, a subset of the Ras-related superfamily of small monomeric GTPases. There are four known isoforms: Rab3a, Rab3b, Rab3c, and Rab3d (1, 2). Rab3a and Rab3c are primarily found in neuronal and neuroendocrine cells, whereas Rab3b and Rab3d are predominantly expressed in non-neuronal tissues such as the parotid gland, pancreas, mast cells, and adipose tissue (2, 3).

Rab3a, the most abundant and well-characterized isoform, is highly enriched in synaptic vesicles. It regulates vesicle transport, docking, fusion, and Ca<sup>2+</sup>-dependent neurotransmitter release by cycling between a GDP-bound inactive state and a GTP-bound vesicle-associated active state, interacting with other synaptic proteins in the process (1, 2).

Unlike integral membrane proteins of synaptic vesicles, Rab3a and Rab3c are absent from the Golgi complex, preventing immunostaining of the axo-dendritic region, which can occur with proteins such as synaptophysin, synaptobrevin/VAMP, or synaptogyrin (1).

Beyond the nervous system, Rab3a is also expressed in the pancreas, where it is predominantly localized to β-cells of the islets of Langerhans. It plays a crucial role in regulated insulin secretion, while Rab3d is primarily involved in exocrine pancreatic enzyme secretion (3).

## Selected References for 107 011

Rab3 proteins involved in vesicle biogenesis and priming in embryonic mouse chromaffin cells.  
Schonn JS, van Weering JR, Mohrmann R, Schlüter OM, Südhof TC, de Wit H, Verhage M, Sørensen JB  
Traffic (Copenhagen, Denmark) (2010) 111: 1415-28. **WB, ICC**

Synaptic targeting of rabphilin-3A, a synaptic vesicle Ca<sup>2+</sup>/phospholipid-binding protein, depends on rab3A/3C.  
Li C, Takei K, Geppert M, Daniell L, Stenius K, Chapman ER, Jahn R, De Camilli P, Südhof TC  
Neuron (1994) 134: 885-98. **WB, IHC**

JIP3 localises to exocytic vesicles and focal adhesions in the growth cones of differentiated PC12 cells.  
Caswell PT, Dickens M  
Molecular and cellular biochemistry (2017) : . **WB, ICC; tested species: rat**

Association of Rab3A with synaptic vesicles at late stages of the secretory pathway.  
Matteoli M, Takei K, Cameron R, Hurlbut P, Johnston PA, Südhof TC, Jahn R, De Camilli P  
The Journal of cell biology (1991) 1153: 625-33. **ICC, WB; tested species: rat**

Microtubule-dependent transport of secretory vesicles in RBL-2H3 cells.  
Smith AJ, Pfeiffer JR, Zhang J, Martinez AM, Griffiths GM, Wilson BS  
Traffic (Copenhagen, Denmark) (2003) 45: 302-12. **EM; tested species: rat**

Monitoring of activity-driven trafficking of endogenous synaptic proteins through proximity labeling.  
Pascual-Caro C, de Juan-Sanz J  
PLoS biology (2024) 2210: e3002860. **WB; tested species: rat**

Kif1a and intact microtubules maintain synaptic-vesicle populations at ribbon synapses in zebrafish hair cells.  
David S, Pinter K, Nguyen KK, Lee DS, Lei Z, Sokolova Y, Sheets L, Kindt KS  
The Journal of physiology (2024) : . **IHC; tested species: zebrafish**

cAMP-EPAC-PKCε-RIM1α signaling regulates presynaptic long-term potentiation and motor learning.  
Wang XT, Zhou L, Dong BB, Xu FX, Wang DJ, Shen EW, Cai XY, Wang Y, Wang N, Ji SJ, Chen W, et al.  
eLife (2023) 12: . **WB; tested species: mouse**

A presynaptic phosphosignaling hub for lasting homeostatic plasticity.  
Müller JA, Betzin J, Santos-Tejedor J, Mayer A, Oprişoreanu AM, Engholm-Keller K, Paulußen I, Gulakova P, McGovern TD, Gschossman LJ, Schönhense E, et al.  
Cell reports (2022) 393: 110696. **WB; tested species: mouse**

Access the online factsheet including applicable protocols  
at <https://sysy.com/product/107011> or scan the QR-code.



# FAQ - How should I store my antibody?

## Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

## Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

## Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

## Product Specific Hints for Storage

### Control proteins / peptides

- Store at -20°C to -80°C.

### Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

### Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

### Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

# Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.