SYSY Antibodies

Rudolf-Wissell-Str. 28a
37079 Göttingen, Germany
Phone: +49 551-50556-0
Fax: +49 551-50556-384
E-mail: sales@sysy.com
Web: www.sysy.com

Syntaxin5

Cat.No. 110 053BT; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen, labeled with Biotin. Albumin was added for stabilization. For **reconstitution** add 50 µl H_2O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	**WB**: 1 : 1000 (AP staining) **IP**: yes **ICC**: 1 : 100 up to 1 : 500 **IHC**: not tested yet **IHC-P (FFPE)**: not tested yet
Label	biotin
Immunogen	Recombinant protein corresponding to AA 1 to 279 from mouse Syntaxin5 (UniProt Id: Q8K1E0)
Reactivity	Reacts with: human (Q13190), rat (Q08851), mouse (Q8K1E0), hamster. Other species not tested yet.
Matching control	110-5P
Remarks	This antibody detects the 35 kDa and 42 kDa variants of syntaxin 5.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Syntaxin 5, a member of the SNARE family of proteins, is functionally related to the yeast protein Sed5p. Two syntaxin 5 isoforms (35 and 42 kDa) have been described which are generated from the same mRNA by alternative translation initiation. The longer 42 kDa variant contains an N-terminal extension carrying a putative type II ER-retrieval signal.
Syntaxin 5 forms a SNARE complex with membrin, GOSR 1, rbet1 and rsec22. A more detailed analysis revealed two subcomplexes within this complex. One contains syntaxin 5 (mainly the shorter 35 kDa variant) and GOSR 1 whereas the other is composed of syntaxin 5 (35 and 42 kDa variant), membrin, rsec22 and rbet1.
Recently syntaxin 5 has been shown to be involved in the processing and accumulation of β-APP in neuronal cells.

Selected General References

Syntaxin 5 interacts specifically with presenilin holoproteins and affects processing of betaAPP in neuronal cells.
Suga K et al. J. Neurochem. (2005) PubMed:15998293

Participation of the syntaxin 5/Ykt6/GS28/GS15 SNARE complex in transport from the early/recycling endosome to the trans-Golgi network.
Tai G et al. Mol. Biol. Cell (2004) PubMed:15215310

rsly1 binding to syntaxin 5 is required for endoplasmic reticulum-to-Golgi transport but does not promote SNARE motif accessibility.
Williams AL et al. Mol. Biol. Cell (2004) PubMed:14565970

GS15 forms a SNARE complex with syntaxin 5, GS28, and Ykt6 and is implicated in traffic in the early cisternae of the Golgi apparatus.
Xu Y et al. Mol. Biol. Cell (2002) PubMed:12388752

Ykt6 forms a SNARE complex with syntaxin 5, GS28, and Bet1 and participates in a late stage in endoplasmic reticulum-Golgi transport.
Zhang T et al. J. Biol. Chem. (2001) PubMed:11323436

Syntaxin 5 is a common component of the NSF- and p97-mediated reassembly pathways of Golgi cisternae from mitotic Golgi fragments in vitro.
Rabouille C et al. Cell (1998) PubMed:9506515

Role of vesicle-associated syntaxin 5 in the assembly of pre-Golgi intermediates.
Rowe T et al. Science (1998) PubMed:9445473

An isoform of the Golgi t-SNARE, syntaxin 5, with an endoplasmic reticulum retrieval signal.
Hui N et al. Mol. Biol. Cell (1997) PubMed:9307973

Cloning and identification of human syntaxin 5 as a synaptobrevin/VAMP binding protein.
Ravichandran V et al. J. Mol. Neurosci. (1997) PubMed:9188044

Mammalian Sly1 regulates syntaxin 5 function in endoplasmic reticulum to Golgi transport.
Dascher C et al. J. Biol. Chem. (1996) PubMed:8663406

Syntaxin 5 regulates endoplasmic reticulum to Golgi transport.
Dascher C et al. J. Biol. Chem. (1994) PubMed:7961911

Access the online factsheet including applicable protocols at https://sysy.com/product/110053BT or scan the QR-code.

FAQ - How should I store my antibody?

Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies**. Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free ("no-frost") freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at –20°C to –80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C

Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C
  Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C
  Adding a carrier protein (e.g., BSA) enhances long-term stability.
  Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C
  Adding a carrier protein such as BSA improves long-term stability.
  Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at –20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.