SYSY Antibodies

Rudolf-Wissell-Str. 28a
37079 Göttingen, Germany
Phone: +49 551-50556-0
Fax: +49 551-50556-384
E-mail: sales@sysy.com
Web: www.sysy.com

Munc13-1

Cat.No. 126 111; Monoclonal mouse antibody, 100 µg purified IgG (lyophilized)

Data Sheet

Reconstitution/ Storage	100 µg purified IgG, lyophilized. Albumin and azide were added for stabilization. For **reconstitution** add 100 µl H_2O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	**WB**: 1 : 500 up to 1 : 5000 (AP staining) **IP**: yes (see remarks) **ICC**: not tested yet **IHC**: not tested yet **IHC-P (FFPE)**: not tested yet **ELISA**:
Clone	266B1
Subtype	IgG2b (κ light chain)
Immunogen	Recombinant protein corresponding to AA 3 to 317 from rat Munc13-1 (UniProt Id: Q62768)
Reactivity	Reacts with: rat (Q62768), mouse (Q4KUS2), zebrafish. Other species not tested yet.
Specificity	K.O. validated
Remarks	**IP**: Denaturing IP-protocol is recommended. Protein-protein interactions may be affected.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Munc13s are homologues of the C. elegans unc-13 gene product. Three brain-specific isoforms, Munc13-1, -13-2, and -13-3, are expressed in mammals, where they localize to presynaptic terminals. All three isoforms share multiple regulatory domains that may mediate phorbol ester and diacylglycerol binding.
Munc13-1 shows the broadest expression pattern and is found in cortex, cerebellum, olfactory bulb, and hippocampus. **Munc13-2** is mainly expressed in cortex and hippocampus, whereas **Munc13-3** exhibits highest expression levels in cerebellum and pons.
Munc13-1 interacts directly with a putative coiled-coil domain in the N-terminal part of syntaxin and is involved in synaptic vesicle priming. For **Munc13-2** an additional ubiquitously expressed N-terminal splice variant (ubMunc13-2) has been described. **Munc13-3** has been shown to be involved in the regulation of cerebellar synaptic transmission and motor learning.

For more information on protein expression pattern, please refer to the overview image in our SYSY Antibodies ATLAS.

Selected References for 126 111

Formation of Golgi-derived active zone precursor vesicles.
Maas C, Torres VI, Altrock WD, Leal-Ortiz S, Wagh D, Terry-Lorenzo RT, Fejtova A, Gundelfinger ED, Ziv NE, Garner CC
The Journal of neuroscience : the official journal of the Society for Neuroscience (2012) 3232: 11095-108. . **WB, ICC**

Different Munc13 isoforms function as priming factors in lytic granule release from murine cytotoxic T lymphocytes.
Dudenhöffer-Pfeifer M, Schirra C, Pattu V, Halimani M, Maier-Peuschel M, Marshall MR, Matti U, Becherer U, Dirks J, Jung M, Lipp P, et al.
Traffic (Copenhagen, Denmark) (2013) 147: 798-809. . **IP; tested species: mouse**

A presynaptic phosphosignaling hub for lasting homeostatic plasticity.
Müller JA, Betzin J, Santos-Tejedor J, Mayer A, Oprişoreanu AM, Engholm-Keller K, Paulußen I, Gulakova P, McGovern TD, Gschossman LJ, Schönhense E, et al.
Cell reports (2022) 393: 110696. . **WB; tested species: mouse**

CYP46A1 Activation by Efavirenz Leads to Behavioral Improvement without Significant Changes in Amyloid Plaque Load in the Brain of 5XFAD Mice.
Petrov AM, Lam M, Mast N, Moon J, Li Y, Maxfield E, Pikuleva IA
Neurotherapeutics : the journal of the American Society for Experimental NeuroTherapeutics (2019) : . . **WB; tested species: mouse**

RIM C2B Domains Target Presynaptic Active Zone Functions to PIP2-Containing Membranes.
de Jong APH, Roggero CM, Ho MR, Wong MY, Brautigam CA, Rizo J, Kaeser PS
Neuron (2018) 982: 335-349.e7. . **ICC; tested species: mouse**

Vti1a/b regulate synaptic vesicle and dense core vesicle secretion via protein sorting at the Golgi.
Emperador-Melero J, Huson V, van Weering J, Bollmann C, Fischer von Mollard G, Toonen RF, Verhage M
Nature communications (2018) 91: 3421. . **ICC; tested species: mouse**

RIM-binding protein 2 regulates release probability by fine-tuning calcium channel localization at murine hippocampal synapses.
Grauel MK, Maglione M, Reddy-Alla S, Willmes CG, Brockmann MM, Trimbuch T, Rosenmund T, Pangalos M, Vardar G, Stumpf A, Walter AM, et al.
Proceedings of the National Academy of Sciences of the United States of America (2016) 11341: 11615-11620. . **WB; tested species: mouse**

Synaptotagmin-12 phosphorylation by cAMP-dependent protein kinase is essential for hippocampal mossy fiber LTP.
Kaeser-Woo YJ, Younts TJ, Yang X, Zhou P, Wu D, Castillo PE, Südhof TC
The Journal of neuroscience : the official journal of the Society for Neuroscience (2013) 3323: 9769-80. . **WB**

Access the online factsheet including applicable protocols at https://sysy.com/product/126111 or scan the QR-code.

FAQ - How should I store my antibody?

Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies**. Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free ("no-frost") freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at –20°C to –80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C

Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C
  Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C
  Adding a carrier protein (e.g., BSA) enhances long-term stability.
  Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C
  Adding a carrier protein such as BSA improves long-term stability.
  Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at –20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.