

## VGLUT1 (SLC17A7)

Cat.No. 135 011; Monoclonal mouse antibody, 100 µg purified IgG (lyophilized)

### Data Sheet

Reconstitution/ Storage	100 µg purified IgG, lyophilized. Albumin and azide were added for stabilization. For <b>reconstitution</b> add 100 µl H <sub>2</sub> O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	<b>WB:</b> 1 : 1000 up to 1 : 5000 (AP staining) (see remarks) <b>IP:</b> yes <b>ICC:</b> 1 : 500 up to 1 : 1000 (see remarks) <b>IHC:</b> 1 : 500 up to 1 : 1000 <b>IHC-P:</b> 1 : 1000
Clone	68B7
Subtype	IgG1 (κ light chain)
Immunogen	Synthetic peptide corresponding to residues near the carboxy terminus of rat VGLUT 1 (UniProt Id: Q62634)
Reactivity	Reacts with: rat (Q62634), mouse (Q3TXX4). Other species not tested yet.
Specificity	K.O. validated
Matching control	135-0P
Remarks	<b>WB:</b> To avoid protein aggregation, do not heat samples for SDS-PAGE. This antibody is highly recommended for Western blot experiments. <b>ICC:</b> The following fixatives are possible: 4% formaldehyde/PFA, acetone.

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

### Background

The vesicular **glutamate transporter 1 VGLUT1**, also referred to as **BNPI** and **SLC17A7**, was originally identified as a brain specific phosphate transporter. Like the related VGLUT2, VGLUT1 is both necessary and sufficient for uptake and storage of glutamate and thus comprises the sole determinant for a glutamatergic phenotype. Both VGLUTs are different from the plasma membrane transporters in that they are driven by a proton electrochemical gradient across the vesicle membrane.

VGLUT1 and VGLUT2 show complementary expression patterns. Together, they are currently the best markers for glutamatergic nerve terminals and glutamatergic synapses.

### Selected References for 135 011

Colocalization of different neurotransmitter transporters on synaptic vesicles is sparse except for VGLUT1 and ZnT3. Upmanyu N, Jin J, Emde HV, Ganzella M, Bösch L, Malviya VN, Zhuleku E, Politi AZ, Ninov M, Silbern I, Leutenegger M, et al. Neuron (2022) : . . **WB, UPTAKE; tested species: rat**

The impact of exogenous Oxytocin on visual cortex plasticity across different stages of visual development. Sun Y, Wang X, Chen Y, Luan Z, Hao R

Scientific reports (2025) 151: 12137. . **WB, IHC; tested species: mouse**

Spatial proteomics in neurons at single-protein resolution.

Unterauer EM, Shetab Boushehri S, Jevdokimenko K, Masullo LA, Ganji M, Sograte-Idrissi S, Kowalewski R, Strauss S, Reinhardt SCM, Perovic A, Marr C, et al.

Cell (2024) 1877: 1785-1800.e16. . **DNA\_PAINT; tested species: rat**

Analyzing schizophrenia-related phenotypes in mice caused by autoantibodies against NRXN1α in schizophrenia.

Shiwaku H, Katayama S, Gao M, Kondo K, Nakano Y, Motokawa Y, Toyoda S, Yoshida F, Hori H, Kubota T, Ishikawa K, et al. Brain, behavior, and immunity (2023) 111: 32-45. . **IHC-P; tested species: mouse**

Neurocalcin Delta Knockout Impairs Adult Neurogenesis Whereas Half Reduction Is Not Pathological.

Upadhyay A, Hosseinibarkoie S, Schneider S, Kaczmarek A, Torres-Benito L, Mendoza-Ferreira N, Overhoff M, Rombo R, Grysko V, Kye MJ, Kononenko NL, et al.

Frontiers in molecular neuroscience (2019) 12: 19. . **ICC; tested species: mouse**

Decreased excitatory and increased inhibitory transmission in the hippocampal CA1 drive neuroinflammation-induced cognitive impairments in mice.

Wu XM, Shi CN, Liu K, Hu XY, He QL, Yao H, Fan D, Ma DQ, Yang JJ, Shen JC, Ji MH, et al.

Brain, behavior, and immunity (2025) 128: 416-428. . **IHC; tested species: mouse**

Neuroimmune Microenvironment Reprogramming via Immuno-piezoelectric Transducers for Synergistic Stem Cell Therapy in Traumatic Brain Injury.

Liang L, Li X, Hu K, Cai P, Wang J, Yu J, Wang S, Zhao Y, Xu C, Li S, Liu H, et al.

Advanced materials (Deerfield Beach, Fla.) (2025) : e12810. . **IHC; tested species: rat**

Environmental enrichment highlights mitochondrial inner membrane function as a therapeutic target for sepsis-associated encephalopathy.

Wang SX, Yin XY, Jian JX, Ji MH, Yang JJ, Shen JC

Brain research (2025) 1864: 149790. . **IHC; tested species: mouse**

FUT8 Regulates Cerebellar Neurogenesis and Development Through Maintaining the Level of Neural Cell Adhesion Molecule Cntn2.

Wei K, Zhang J, Qu W, Zhu J, Zhu Q, Yi W, Zou C, Ma D, Li X

Molecular neurobiology (2025) 625: 5679-5694. . **IHC; tested species: mouse**

Low-dose cannabidiol treatment prevents chronic stress-induced phenotypes and is associated with multiple synaptic changes across various brain regions.

Borràs-Pernas S, Sancho-Balsells A, Patterer L, Wang M, Del Toro D, Alberch J, Schibano D, Espel J, Heybeck M, Scheidel B, Giral A, et al.

Neuropharmacology (2025) 277: 110526. . **IHC; tested species: mouse**

Access the online factsheet including applicable protocols at <https://sysy.com/product/135011> or scan the QR-code.



## FAQ - How should I store my antibody?

### Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

### Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

### Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

### Product Specific Hints for Storage

#### Control proteins / peptides

- Store at -20°C to -80°C.

#### Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

#### Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

#### Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

## Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.