

VGLUT1

Cat.No. 135 308; Recombinant rabbit antibody, 50 µg recombinant IgG (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg purified recombinant IgG, lyophilized. Albumin and azide were added for stabilization. For reconstitution add 50 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: 1 : 1000 up to 1 : 5000 (AP staining) (see remarks) IP: yes ICC: 1 : 1000 IHC: 1 : 500 up to 1 : 1000 IHC_P: 1 : 1000 EXM: yes (see remarks)
Clone	Rb68B7
Subtype	IgG1 (κ light chain)
Immunogen	Synthetic peptide corresponding to residues near the carboxy terminus of rat VGLUT 1. (UniProt Id: Q62634)
Reactivity	Reacts with: rat (Q62634), mouse (Q3TXX4). Other species not tested yet.
Specificity	K.O.
Matching control	135-0P
Remarks	This antibody is a chimeric antibody based on the monoclonal mouse antibody clone 68B7. The constant regions of the heavy and light chains have been replaced by rabbit specific sequences. Therefore, the antibody can be used with standard anti-rabbit secondary reagents. The antibody has been expressed in mammalian cells. WB: This antibody is highly recommended for Western blot experiments. VGLUT 1 aggregates after boiling, making it necessary to run SDS-PAGE with non-boiled samples. EXM: This antibody has been successfully used for the epitope-preserving magnified analysis of the proteome (eMAP) expansion microscopy method (Park et al. 2021. PMID: 34767453).

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

The vesicular **glutamate transporter 1 VGLUT 1**, also referred to as **BNPI** and **SLC17A7**, was originally identified as a brain specific phosphate transporter. Like the related VGLUT 2, VGLUT 1 is both necessary and sufficient for uptake and storage of glutamate and thus comprises the sole determinant for a glutamatergic phenotype. Both VGLUTs are different from the plasma membrane transporters in that they are driven by a proton electrochemical gradient across the vesicle membrane.

VGLUT 1 and VGLUT 2 show complementary expression patterns. Together, they are currently the best markers for glutamatergic nerve terminals and glutamatergic synapses.

Selected References for 135 308

- Dorsal and ventral striatal neuronal subpopulations differentially disrupt male mouse copulatory behavior. Detraux B, Vilella A, De Groote A, Schiffmann SN, Zoli M, de Kerchove d'Exaerde A. *European neuropsychopharmacology : the journal of the European College of Neuropsychopharmacology* (2021) 49: 23-37. . **IHC; tested species: mouse**
- Experimental Epileptogenesis in a Cell Culture Model of Primary Neurons from Rat Brain: A Temporal Multi-Scale Study. Jablonski J, Hoffmann L, Blümcke I, Fejtová A, Uebe S, Ekici AB, Gnatkovsky V, Kobow K. *Cells* (2021) 1011: . . **ICC; tested species: rat**
- Tonic activin signaling shapes cellular and synaptic properties of CA1 neurons mainly in dorsal hippocampus. Dahlmans M, Valero-Aracama MJ, Dahlmans JK, Zheng F, Alzheimer C. *iScience* (2023) 2610: 108001. . **ICC; tested species: mouse**
- Glutamate released by Cajal-Retzius cells impacts specific hippocampal circuits and behaviors. Anstötz M, Lee SK, Maccaferri G. *Cell reports* (2022) 397: 110822. . **IHC; tested species: mouse**
- Melatonin pretreatment alleviates the long-term synaptic toxicity and dysmyelination induced by neonatal Sevoflurane exposure via MT1 receptor-mediated Wnt signaling modulation. Liang L, Zeng T, Zhao Y, Lu R, Guo B, Xie R, Tang W, Zhang L, Mao Z, Yang X, Wu S, et al. *Journal of pineal research* (2021) 714: e12771. . **IHC; tested species: mouse**

Selected General References

- Identification of a vesicular glutamate transporter that defines a glutamatergic phenotype in neurons. Takamori S, Rhee JS, Rosenmund C, Jahn R. *Nature* (2000) 4076801: 189-94. .
- Uptake of glutamate into synaptic vesicles by an inorganic phosphate transporter. Bellocchio EE, Reimer RJ, Fremerey RT, Edwards RH. *Science (New York, N.Y.)* (2000) 2895481: 957-60. .
- The localization of the brain-specific inorganic phosphate transporter suggests a specific presynaptic role in glutamatergic transmission. Bellocchio EE, Hu H, Pohorille A, Chan J, Pickel VM, Edwards RH. *The Journal of neuroscience : the official journal of the Society for Neuroscience* (1998) 1821: 8648-59. .
- Cloning and expression of a cDNA encoding a brain-specific Na(+)-dependent inorganic phosphate cotransporter. Ni B, Rosteck PR, Nadi NS, Paul SM. *Proceedings of the National Academy of Sciences of the United States of America* (1994) 9112: 5607-11. .

Access the online factsheet including applicable protocols at <https://sysy.com/product/135308> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.