

Ca²⁺ channel P/Q-type α -1A

Cat.No. 152 211; Monoclonal mouse antibody, 100 μ g purified IgG (lyophilized)

Data Sheet

Reconstitution/ Storage	100 μ g purified IgG, lyophilized. Azide was added before lyophilization. For reconstitution add 100 μ l H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: 1 : 1000 (AP staining) (see remarks) IP: not tested yet ICC: 1 : 100 up to 1 : 500 (see remarks) IHC: not tested yet IHC-P: not tested yet IHC-Fr: 1 : 500 (see remarks)
Clone	23F11
Subtype	IgG2b (κ light chain)
Immunogen	Recombinant protein corresponding to AA 1921 to 2212 from rat Ca ²⁺ channel P/Q-type α -1A (Cav2.1) (UniProt Id: P54282)
Reactivity	Reacts with: rat (P54282), mouse (P97445). Other species not tested yet.
Remarks	WB: Less sensitive than the polyclonal antibodies (152 203, 152 205). Due to its large size, this antibody requires special gel-electrophoresis and Western blot protocols for visualization by immunoblotting. Excellent results can be obtained with the 4-12% TRIS-glycine gradient gels of anamed or NuPage TRIS-acetate gels from Invitrogen. Ca ²⁺ channel P/Q-type α -1A aggregates after boiling, making it necessary to run SDS-PAGE with non-boiled samples. ICC: This antibody requires mild fixation (1-2% PFA for 15min). IHC-Fr: Formaldehyde, acetone, methanol or methanol-acetone fixation is recommended.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Voltage gated calcium channels (VGCCs), also referred to as voltage sensitive calcium channels (VSCCs), are present in most excitable cells. They mediate the influx of Ca²⁺ ions into the cell and trigger the release of neurotransmitters or hormones but are also involved in other calcium dependent processes like metabolism, cell proliferation and cell death.

VGCCs are composed of four subunits (α -1, α -2, β and δ) in a 1:1:1:1 ratio. The α -1A isoform occurs in VGCCs of the **P/Q-type** while isoform α -1B is found in the N-type. Both belong to the high voltage activated group (hva).

Selected References for 152 211

Transient Confinement of CaV2.1 Ca²⁺-Channel Splice Variants Shapes Synaptic Short-Term Plasticity. Heck J, Parutto P, Ciuraszkiewicz A, Bikbaev A, Freund R, Mitlöhner J, Alonso M, Fejtova A, Holcman D, Heine M Neuron (2019) . . . **WB; tested species: rat**

Selected General References

Calcium channel types with distinct presynaptic localization couple differentially to transmitter release in single calyx-type synapses.

Wu LG et al. J. Neurosci. (1999) PubMed:9880593

Localization of Ca²⁺ channel subtypes on rat spinal motor neurons, interneurons, and nerve terminals.

Westenbroek RE et al. J. Neurosci. (1998) PubMed:9698323

Biochemical properties and subcellular distribution of the BI and rBA isoforms of alpha 1A subunits of brain calcium channels.

Sakurai T et al. J. Cell Biol. (1996) PubMed:8707834

Immunochemical identification and subcellular distribution of the alpha 1A subunits of brain calcium channels.

Westenbroek RE et al. J. Neurosci. (1995) PubMed:7472404

Immunochemical identification and differential phosphorylation of alternatively spliced forms of the alpha 1A subunit of brain calcium channels.

Sakurai T et al. J. Biol. Chem. (1995) PubMed:7673157

Primary structure of a calcium channel that is highly expressed in the rat cerebellum.

Starr TV et al. Proc. Natl. Acad. Sci. U.S.A. (1991) PubMed:1648226

Access the online factsheet including applicable protocols at <https://sysy.com/product/152211> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.