

## Veli1/2/3

Cat.No. 184 002; Polyclonal rabbit antibody, 200 µl antiserum (lyophilized)

### Data Sheet

Reconstitution/ Storage	200 µl antiserum, lyophilized. For <b>reconstitution</b> add 200 µl H <sub>2</sub> O, then aliquot and store at -20°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	<b>WB:</b> 1 : 1000 (AP staining) <b>IP:</b> not tested yet <b>ICC:</b> 1 : 1000 <b>IHC:</b> not tested yet <b>IHC-P:</b> not tested yet
Immunogen	Recombinant protein corresponding to AA 1 to 232 from rat Velis1 (UniProt Id: Q9Z250)
Reactivity	Reacts with: human (O14910, Q9HAP6, Q9NUP9), rat (Q9Z250, Q9Z252, Q79210), mouse (Q8JZS0, O88951, O88952). Other species not tested yet.
Specificity	Recognizes all three isoforms veli 1, 2 and 3.
Matching control	184-0P

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

### Background

The complex of the multi-adaptor-proteins CASK, Mint and **Veli**, also referred to as **MALS** and **Lin 7**, is conserved across kingdoms and may be involved in localizing synaptic vesicles at synaptic terminals. Three Veli isoforms (1, 2 and 3) with different expression patterns have been described, so far. Veli proteins have also been suggested to participate in regulating the recruitment of neurotransmitter receptors to the post synaptic density (PSD) by interaction with their PDZ domains.

### Selected References for 184 002

Functional analysis of CASK transcript variants expressed in human brain.  
Tibbe D, Pan YE, Reißner C, Harms FL, Kreienkamp HJ  
PloS one (2021) 166: e0253223. . **WB; tested species: human**

Patient-Derived Variants Define Constraints for Ligand Binding at the PDZ Domain of CASK.  
Tibbe D, Hönick HH, Bhatia N, Truong T, Proskauer L, Ortiz-Gonzalez X, Maguire JA, Pak C, Kreienkamp HJ  
Journal of neurochemistry (2025) 16912: e70303. . **WB; tested species: human**

Liprin-α proteins are master regulators of human presynapse assembly.  
Marcó de la Cruz B, Campos J, Molinaro A, Xie X, Jin G, Wei Z, Acuna C, Sterky FH  
Nature neuroscience (2024) : . . **WB; tested species: human**

Regulation of Liprin-α phase separation by CASK is disrupted by a mutation in its CaM kinase domain.  
Tibbe D, Ferle P, Krisp C, Nampoothiri S, Mirzaa G, Assaf M, Parikh S, Kutsche K, Kreienkamp HJ  
Life science alliance (2022) 510: . . **WB; tested species: human**

### Selected General References

Synaptic transmission regulated by a presynaptic MALS/Liprin-alpha protein complex.  
Olsen O et al. Curr. Opin. Cell Biol. (2006) PubMed:16504495

Neurotransmitter release regulated by a MALS-liprin-alpha presynaptic complex.  
Olsen O et al. J. Cell Biol. (2005) PubMed:16186258

The SNARE motif contributes to rbet1 intracellular targeting and dynamics independently of SNARE interactions.  
Joglekar AP et al. J. Biol. Chem. (2003) PubMed:12566453

Access the online factsheet including applicable protocols at <https://sysy.com/product/184002> or scan the QR-code.



# FAQ - How should I store my antibody?

## Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

## Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies.** Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

## Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free (“no-frost”) freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at -20°C to -80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

## Product Specific Hints for Storage

### Control proteins / peptides

- Store at -20°C to -80°C

### Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C. Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C. Adding a carrier protein (e.g., BSA) enhances long-term stability. Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

### Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C. Adding a carrier protein such as BSA improves long-term stability. Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

### Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

# Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at -20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.