SYSY Antibodies

Rudolf-Wissell-Str. 28a
37079 Göttingen, Germany
Phone: +49 551-50556-0
Fax: +49 551-50556-384
E-mail: sales@sysy.com
Web: www.sysy.com

PACSIN1

Cat.No. 196 002; Polyclonal rabbit antibody, 200 µl antiserum (lyophilized)

Data Sheet

Reconstitution/ Storage	200 µl antiserum, lyophilized. For **reconstitution** add 200 µl H_2O, then aliquot and store at -20°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	**WB**: 1 : 1000 (AP staining) **IP**: not tested yet **ICC**: 1 : 500 up to 1 : 1000 **IHC**: not tested yet **IHC-P (FFPE)**: not tested yet
Immunogen	Recombinant protein corresponding to AA 293 to 384 from mouse PACSIN1 (UniProt Id: Q61644)
Reactivity	Reacts with: rat (Q9Z0W5), mouse (Q61644). Other species not tested yet.
Matching control	196-0P

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

This protein has been described as **PACSIN 1** (for **P**KC **a**nd **C**K2 **s**ubstrate **i**n **n**eurons) and **syndapin 1** (**syn**aptic, **d**ynamin-**a**ssociated **p**rote**in** I). It is a multidomain phospho-protein that interacts via its SH3 domain with dynamin 1, synaptojanin, synapsin and N-WASP. Since N-WASP is an activator of the Arp2/3 complex, syndapin may link membrane trafficking and the actin cytoskeleton.
Up to now, 3 isoforms have been described. Syndapin 1 shows a neuronal, syndapin 3 a muscular and syndapin 2 an ubiquitous expression profile.

Selected References for 196 002

A local, periactive zone endocytic machinery at photoreceptor synapses in close vicinity to synaptic ribbons.
Wahl S, Katiyar R, Schmitz F
The Journal of neuroscience : the official journal of the Society for Neuroscience (2013) 3325: 10278-300. . **WB, ICC**

Composition of isolated synaptic boutons reveals the amounts of vesicle trafficking proteins.
Wilhelm BG, Mandad S, Truckenbrodt S, Kröhnert K, Schäfer C, Rammner B, Koo SJ, Claßen GA, Krauss M, Haucke V, Urlaub H, et al.
Science (New York, N.Y.) (2014) 3446187: 1023-8. . **ICC, IHC; tested species: mouse,rat**

A novel glycine receptor variant with startle disease affects syndapin I and glycinergic inhibition.
Langlhofer G, Schaefer N, Maric HM, Keramidas A, Zhang Y, Baumann P, Blum R, Breitinger U, Strømgaard K, Schlosser A, Kessels MM, et al.
The Journal of neuroscience : the official journal of the Society for Neuroscience (2020) : . . **WB, ICC; tested species: human,mouse**

Native KCC2 interactome reveals PACSIN1 as a critical regulator of synaptic inhibition.
Mahadevan V, Khademullah CS, Dargaei Z, Chevrier J, Uvarov P, Kwan J, Bagshaw RD, Pawson T, Emili A, De Koninck Y, Anggono V, et al.
eLife (2017) 6: . . **IP; tested species: mouse**

Impaired Presynaptic Function Contributes Significantly to the Pathology of Glycine Receptor Autoantibodies.
Wiessler AL, Zheng F, Werner C, Habib M, Tuzun E, Alzheimer C, Sommer C, Villmann C
Neurology(R) neuroimmunology & neuroinflammation (2025) 122: e200364. . **WB; tested species: mouse**

Dynamin is primed at endocytic sites for ultrafast endocytosis.
Imoto Y, Raychaudhuri S, Ma Y, Fenske P, Sandoval E, Itoh K, Blumrich EM, Matsubayashi HT, Mamer L, Zarebidaki F, Söhl-Kielczynski B, et al.
Neuron (2022) 11017: 2815-2835.e13. . **ICC; tested species: mouse**

Synaptic AP2 CCV life cycle regulation by the Eps15, ITSN1, Sgip1/AP2, synaptojanin1 interactome.
Mishra R, Sengül GF, Candiello E, Schu P
Scientific reports (2021) 111: 8007. . **WB; tested species: mouse**

Selected General References

Syndapin I is the phosphorylation-regulated dynamin I partner in synaptic vesicle endocytosis.
Anggono V et al. Nat. Neurosci. (2006) PubMed:16648848

Syndapin oligomers interconnect the machineries for endocytic vesicle formation and actin polymerization.
Kessels MM et al. J. Biol. Chem. (2006) PubMed:16540475

The syndapin protein family: linking membrane trafficking with the cytoskeleton.
Kessels MM et al. J. Cell. Sci. (2004) PubMed:15226389

All three PACSIN isoforms bind to endocytic proteins and inhibit endocytosis.
Modregger J et al. J. Cell. Sci. (2000) PubMed:11082044

Access the online factsheet including applicable protocols at https://sysy.com/product/196002 or scan the QR-code.

FAQ - How should I store my antibody?

Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies**. Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free ("no-frost") freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at –20°C to –80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C

Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C
  Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C
  Adding a carrier protein (e.g., BSA) enhances long-term stability.
  Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C
  Adding a carrier protein such as BSA improves long-term stability.
  Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at –20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.