

## Abeta38/40/42/43

Cat.No. 218 113; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

### Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Albumin was added for stabilization. For <b>reconstitution</b> add 50 µl H <sub>2</sub> O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	<b>WB:</b> not tested yet <b>IP:</b> not tested yet <b>ICC:</b> not tested yet <b>IHC:</b> 1 : 100 up to 1 : 500 <b>IHC-P:</b> 1 : 1000 (see remarks) <b>ELISA:</b> not tested yet
Immunogen	Synthetic peptide corresponding to AA 13 to 29 from human Abeta (UniProt Id: P05067)
Reactivity	Reacts with: human (P05067), mouse (P12023), dog. Other species not tested yet.
Specificity	Recognizes Abeta 38, 40, 42, 43. Due to sequence homology, a cross-reactivity to unprocessed APP cannot be excluded.
Remarks	<b>IHC:</b> Antigen retrieval with formic acid is required. <b>IHC-P:</b> Antigen retrieval with citrate buffer pH 6, followed by formic acid treatment, is required. Antigen retrieval with formic acid increases sensitivity and reveals more plaques.

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

### Background

Amyloid deposits, also called plaques, of Alzheimer's patients consist of several protein components like the amyloid **beta**-peptides (**Abeta**, **Aβ**) 1-40/42/43 and additional C- and N-terminally modified fragments of Abeta as for instance Abeta pE3 and Abeta pE11.

An additional Abeta variant, **Abeta38**, is more soluble compared to other Abeta species and is not found in plaques of sporadic Alzheimer's cases. However, it is detected in the blood-vessel walls of a subset of patients with severe cerebral amyloid angiopathy. It especially accumulates in brains of patients carrying mutations in the Abeta coding region.

Cleavage of amyloid precursor protein APP by β- and γ- secretases results in the generation of the Aβ (BA4)peptide, whereas α-secretase cleaves within the Aβ sequence and prevents the formation of Abeta from APP.

### Selected References for 218 113

Detection and Quantification of β-Amyloid, Pyroglutamate Aβ, and Tau in Aged Canines.  
Schmidt F, Boltze J, Jäger C, Hofmann S, Willems N, Seeger J, Härtig W, Stolzing A  
Journal of neuropathology and experimental neurology (2015) 749: 912-23. . **IHC**

Aβ38 in the brains of patients with sporadic and familial Alzheimer's disease and transgenic mouse models.  
Reinert J, Martens H, Huettnerrauch M, Kolbow T, Lannfelt L, Ingelsson M, Paetau A, Verkoniemi-Ahola A, Bayer TA, Wirths O  
Journal of Alzheimer's disease : JAD (2014) 394: 871-81. . **WB; tested species: human**

Neurovascular Specifications in the Alzheimer-Like Brain of Mice Affected by Focal Cerebral Ischemia: Implications for Future Therapies.

Michalski D, Hofmann S, Pitsch R, Grosche J, Härtig W  
Journal of Alzheimer's disease : JAD (2017) 592: 655-674. . **IHC; tested species: mouse**

### Selected General References

Circulating immune complexes of Abeta and IgM in plasma of patients with Alzheimer's disease.  
Marcello A et al. J Neural Transm (Vienna) (2009) PubMed:19415450

Immune response to Abeta-peptides in peripheral blood from patients with Alzheimer's disease and control subjects.  
Baril L et al. Neurosci. Lett. (2004) PubMed:14732472

Dietary Cu stabilizes brain superoxide dismutase 1 activity and reduces amyloid Abeta production in APP23 transgenic mice.  
Bayer TA et al. Proc. Natl. Acad. Sci. U.S.A. (2003) PubMed:14617773

Correlative memory deficits, Abeta elevation, and amyloid plaques in transgenic mice.  
Hsiao K et al. Science (1996) PubMed:8810256

Physical, morphological and functional differences between ph 5.8 and 7.4 aggregates of the Alzheimer's amyloid peptide Abeta.

Wood SJ et al. J. Mol. Biol. (1996) PubMed:8601838

Water-soluble Abeta (N-40, N-42) oligomers in normal and Alzheimer disease brains.  
Kuo YM et al. J. Biol. Chem. (1996) PubMed:8626743

Access the online factsheet including applicable protocols  
at <https://sysy.com/product/218113> or scan the QR-code.



## FAQ - How should I store my antibody?

### Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

### Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

### Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

### Product Specific Hints for Storage

#### Control proteins / peptides

- Store at -20°C to -80°C.

#### Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

#### Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

#### Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

## Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.