

M6a Glycoprotein

Cat.No. 238 003; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Albumin was added for stabilization. For reconstitution add 50 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: 1 : 1000 (AP staining) IP: yes ICC: 1 : 1000 IHC: 1 : 200 IHC-P: 1 : 200
Immunogen	Synthetic peptide corresponding to AA 261 to 270 from rat M6a Glycoprotein (UniProt Id: Q812E9)
Reactivity	Reacts with: human (P51674), rat (Q812E9), mouse (P35802), pig, dog, zebrafish. Other species not tested yet.
Matching control	238-0P

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

M6a and **M6b** are transmembrane **glycoproteins** with four membrane spanning domains that belong to the myelin proteolipid protein (PLP) family. In contrast to other PLP proteins that are expressed by glia cells, **M6a** is found only in neurons where it is concentrated at presynaptic glutamateric terminals. **M6a** has been suggested to be a potential mediator of cell-cell interactions involved in axonfasciculation during development.

Selected References for 238 003

Serotonin Transporter Associated Protein Complexes Are Enriched in Synaptic Vesicle Proteins and Proteins Involved in Energy Metabolism and Ion Homeostasis.

Haase J, Grudzinska-Goebel J, Müller HK, Münster-Wandowski A, Chow E, Wynne K, Farsi Z, Zander JF, Ahnert-Hilger G ACS chemical neuroscience (2017) 85: 1101-1116. . **WB; tested species: mouse, rat**

Chronic pharmacologic manipulation of dopamine transmission ameliorates metabolic disturbance in trappc9-linked brain developmental syndrome.

Li Y, Muhammad U, Sapp E, Ke Y, Wang Z, Boudi A, DiFiglia M, Li X JCI insight (2024) : . . **WB; tested species: mouse**

Selected General References

Expression of the axonal membrane glycoprotein M6a is regulated by chronic stress.

Cooper B et al. PLoS ONE (2009) PubMed:19180239

Glycoprotein M6a is present in glutamatergic axons in adult rat forebrain and cerebellum.

Cooper B et al. Brain Res. (2008) PubMed:18241840

The stress-regulated protein M6a is a key modulator for neurite outgrowth and filopodium/spine formation.

Alfonso J et al. Proc. Natl. Acad. Sci. U.S.A. (2005) PubMed:16286650

Immunoelectron microscopic localization of the M6a antigen in rat brain.

Roussel G et al. J. Neurocytol. (1998) PubMed:10447243

Molecular cloning of M6: identification of a PLP/DM20 gene family.

Yan Y et al. Neuron (1993) PubMed:8398137

EMA: a developmentally regulated cell-surface glycoprotein of CNS neurons that is concentrated at the leading edge of growth cones.

Baumrind NL et al. Dev. Dyn. (1992) PubMed:1286213

Access the online factsheet including applicable protocols at <https://sysy.com/product/238003> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies.** Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free (“no-frost”) freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at -20°C to -80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C

Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C. Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C. Adding a carrier protein (e.g., BSA) enhances long-term stability. Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C. Adding a carrier protein such as BSA improves long-term stability. Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at -20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.