

Numblike

Cat.No. 374-0P; control peptide, 100 µg peptide (lyophilized)

Data Sheet

Rudolf-Wissell-Str. 28a

Phone:

E-mail:

Web:

Fax:

37079 Göttingen, Germany

+49 551-50556-0

sales@sysy.com

www.sysy.com

+49 551-50556-384

Reconstitution/ Storage	100 μg peptide, lyophilized. For reconstitution add 100 μl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Control peptides should be stored at -20°C when still lyophilized! For detailed information, see back of the data sheet.
Immunogen	Synthetic peptide corresponding to AA 579 to 595 from mouse Numbl (UniProt Id: O08919)
Recommended dilution	Optimal concentrations should be determined by the end-user.
Matching antibodies	374 003
Remarks	This control peptide consists of the synthetic peptide (aa 579-595 of mouse Numblike) that has been used for immunization. It has been tested in preadsorption experiments and blocks efficiently and specifically the corresponding signal in Western blots. The amount of peptide needed for efficient blocking depends on the titer and on the affinity of the antibody to the antigen.

TO BE USED IN VITRO / FOR RESEARCH ONLY NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Numb proteins (**Numblike** and Numb) display a complex pattern of functions such as the control of asymmetric cell division, cell fate choice, endocytosis, cell adhesion, and cell migration. They have been shown to inhibit Notch signaling by stimulating endocytosis of Notch.

Numb and Numblike have at least partially distinct functions. Numblike is a negative regulator of the NF-KB signaling pathway by abrogating TRAF5-induced activation of NF-KB. Recently, Numblike was implicated as a physiologically relevant target of microRNA miR-34a in neural progenitor cells allowing for enhanced Notch signaling and inhibition of neuronal differentiation. Numb and Numblike are essential in maintaining neural progenitor cells during early neurogenesis by allowing cells to choose progenitor over neuronal fates. They were recently also discovered to be involved in cardiac morphogenesis.

Selected General References

Numb family proteins: novel players in cardiac morphogenesis and cardiac progenitor cell differentiation. Wu M et al. Biomol Concepts (2015) PubMed:25883210

Precardiac deletion of Numb and Numblike reveals renewal of cardiac progenitors. Shenje LT et al. Elife (2014) PubMed:24843018

Numblike regulates proliferation, apoptosis, and invasion of lung cancer cell. Yingjie L et al. Tumour Biol. (2013) PubMed:23681800

Inhibition of Notch2 by Numb/Numblike controls myocardial compaction in the heart. Yang J et al. Cardiovasc. Res. (2012) PubMed:22865640

MiR-34a represses Numbl in murine neural progenitor cells and antagonizes neuronal differentiation. Fineberg SK et al. PLoS ONE (2012) PubMed:22701667

Numbl inhibits glioma cell migration and invasion by suppressing TRAF5-mediated NF-κB activation. Tao T et al. Mol. Biol. Cell (2012) PubMed:22593207

Numblike and Numb differentially affect p53 and Sonic Hedgehog signaling. Liu L et al. Biochem. Biophys. Res. Commun. (2011) PubMed:21893032

The multiple functions of Numb. Gulino A et al. Exp. Cell Res. (2010) PubMed:19944684

Numb and Numbl are required for maintenance of cadherin-based adhesion and polarity of neural progenitors. Rasin MR et al. Nat. Neurosci. (2007) PubMed:17589506

Postnatal deletion of Numb/Numblike reveals repair and remodeling capacity in the subventricular neurogenic niche. Kuo CT et al. Cell (2006) PubMed:17174898

Continuing role for mouse Numb and Numbl in maintaining progenitor cells during cortical neurogenesis. Petersen PH et al. Nat. Neurosci. (2004) PubMed:15273690

Differential expression of mammalian Numb, Numblike and Notch1 suggests distinct roles during mouse cortical neurogenesis. Zhong W et al. Development (1997) PubMed:9169836

Access the online factsheet including applicable protocols at <u>https://sysy.com/product/374-0P</u> or scan the QR-code.

FAQ - How should I store my antibody?

Shipping Conditions

• All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freezedried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- Unlabeled and biotin-labeled antibodies and control proteins should be stored at 4°C before reconstitution. They must not be stored in the freezer when still lyophilized! Temperatures below zero may cause loss of performance.
- Fluorescence-labeled antibodies should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 μl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

• Store at -20°C to -80°C.

Monoclonal Antibodies

- Ascites and hybridoma supernatant should be stored at -20°C up to -80°C. Prolonged storage at 4°C is not recommended! Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera**: With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- Affinity purified antibodies: Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

• Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freezethaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.