

Brn3a

Cat.No. 411 003; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

Data Sheet

Reconstitution/	50 µg specific antibody, lyophilized. Affinity purified with the immunogen.
Storage	Albumin and azide were added for stabilization. For reconstitution add 50 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: not tested yet IP: not tested yet ICC: not tested yet IHC: 1 : 1000 up to 1 : 5000 (see remarks) IHC-P: 1 : 500 up to 1 : 1000
Immunogen	Synthetic peptide corresponding to residues near the amino terminus of mouse Brn3a (UniProt Id: P17208)
Reactivity	Reacts with: mouse (P17208). Other species not tested yet.
Remarks	IHC: For optimal results in retina tissue, follow the retina protocol.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Brn3a, also referred to as POU4F1, RGC-1 or Oct-T1, is a transcription factor highly expressed in the developing peripheral sensory nervous system, in cells of the B- and T-lymphocytic lineages and in certain regions of the CNS e.g. retina, spinal cord, midbrain superior colliculus, red nucleus, nucleus ambiguus, inferior olfactory nucleus and habenula. In the retina Brn3a is a well-established marker for retinal ganglion cells.

Selected References for 411 003

Resveratrol attenuates retinal ganglion cell loss in a mouse model of retinal ischemia reperfusion injury via multiple pathways. Ji K, Li Z, Lei Y, Xu W, Ouyang L, He T, Xing Y. *Experimental eye research* (2021) 209: 108683. . **WB, IHC; tested species: mouse**

Selected Ionotropic Receptors and Voltage-Gated Ion Channels: More Functional Competence for Human Induced Pluripotent Stem Cell (iPSC)-Derived Nociceptors. Schoepf CL, Zeidler M, Specker L, Kern G, Lechner J, Kummer KK, Kress M. *Brain sciences* (2020) 106: . . **ICC; tested species: human, mouse**

Retinal ganglion cell survival and functional maturation in transiently vascularized human retinal organoids. Sharma K, Habibey R, Ribeiro MM, Cui B, Siwicki RA, Striebel J, Pawlick JS, Zorn J, Utz L, Renner M, Picelli S, et al. *Cell stem cell* (2026) : . . **ICC; tested species: human**

An improved method of transducing retinal ganglion cells using AAV via transpupillary injection in adult mouse eyes. Lin F, Lin ST, Wang J, Sellers JT, Chrenek MA, Nickerson JM, Boatright JH, Geisert EE. *Molecular vision* (2025) 31: 1-9. . **IHC; tested species: mouse**

Differential Expression of the Synapse Regulatory Proteins Neurexins in Early Diabetic Retinal Disease. Unlu EK, Marx-Rattner R, Klein KA, Dawson VL, Dawson TM, Sachdeva MM. *Investigative ophthalmology & visual science* (2025) 6611: 46. . **ICC; tested species: human**

The neuronal chaperone proSAAS is highly expressed in the retina. Schaffer N, Mitiás S, Guo Y, Bernstein SL, Lindberg I. *PloS one* (2025) 205: e0321867. . **IHC; tested species: mouse**

Myeloid lineage C3 induces reactive gliosis and neuronal stress during CNS inflammation. Garton T, Smith MD, Kesharwani A, Gharagozloo M, Oh S, Na CH, Absinta M, Reich DS, Zack DJ, Calabresi PA. *Nature communications* (2025) 161: 3481. . **IHC; tested species: mouse**

Mettl3-Mediated N6-Methyladenosine Modification Mitigates Ganglion Cell Loss and Retinal Dysfunction in Retinal Ischemia-Reperfusion Injury by Inhibiting FoxO1-Mediated Autophagy. Zhu F, Feng J, Pan Y, Ouyang L, He T, Xing Y. *Investigative ophthalmology & visual science* (2025) 662: 58. . **IHC; tested species: mouse**

NLRX1 limits inflammatory neurodegeneration in the anterior visual pathway. Gill AJ, Smith MD, Galleguillos D, Garton T, Mace JW, Gadani SP, Kumar S, Pokharel A, Solem K, Potluri S, Hussein O, et al. *Journal of neuroinflammation* (2025) 221: 21. . **IHC; tested species: mouse**

Ameliorative effect of resveratrol on acute ocular hypertension induced retinal injury through the SIRT1/NF-κB pathway. Ji KB, Wan W, Yang Y, He XJ, Xing YQ, Hu Z. *Neuroscience letters* (2024) 826: 137712. . **IHC; tested species: mouse**

AAV2 vector optimization for retinal ganglion cell-targeted delivery of therapeutic genes. Chaour B, Duong TT, Yue J, Liu T, Camacho D, Dine KE, Esteve-Rudd J, Ellis S, Bennett J, Shindler KS, Ross AG, et al. *Gene therapy* (2024) 313-4: 175-186. . **IHC; tested species: mouse**

Optimizing retinal ganglion cell nuclear staining for automated cell counting. Lin F, Lin ST, Wang J, Geisert EE. *Experimental eye research* (2024) 242: 109881. . **IHC; tested species: mouse**

Access the online factsheet including applicable protocols at <https://sysy.com/product/411003> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled and biotin-labeled antibodies and control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites and hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.