

CPT1c

Cat.No. 415 003; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Azide was added before lyophilization. For reconstitution add 50 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: 1 : 1000 (AP staining) IP: yes ICC: 1 : 500 IHC: 1 : 500 up to 1 : 1000 IHC-P: not tested yet
Immunogen	Synthetic peptide corresponding to AA 784 to 798 from mouse Cpt1c (UniProt Id: Q8BGD5)
Reactivity	Reacts with: mouse (Q8BGD5), rat (F1LN46). Other species not tested yet.
Specificity	K.O. validated PubMed: 37309891
Matching control	415-0P

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

CPT1c is a carnitine palmitoyltransferase 1 (CPT 1) isoform that is expressed only in the brain. It has been shown to be an integral part of native AMPA receptor complexes in neurons and modulates the surface expression of these ionotropic glutamate receptors.


Selected References for 415 003

CPT1C is required for synaptic plasticity and oscillatory activity that supports motor, associative and non-associative learning. Iborra-Lázaro G, Djebari S, Sánchez-Rodríguez I, Gratacós-Batlle E, Sánchez-Fernández N, Radošević M, Casals N, Navarro-López JD, Soto Del Cerro D, Jiménez-Díaz L. The Journal of physiology (2023) : . IHC; KO verified; tested species: mouse

Selected General References

Mechanisms of CPT1C-Dependent AMPAR Trafficking Enhancement. Gratacós-Batlle E et al. Front Mol Neurosci (2018) PubMed:30135643
Cpt1c regulated by AMPK promotes papillary thyroid carcinomas cells survival under metabolic stress conditions. Wang R et al. J Cancer (2017) PubMed:29151954
Novel Regulation of the Synthesis of α-Amino-3-hydroxy-5-methyl-4-isoxazolepropionic Acid (AMPA) Receptor Subunit GluA1 by Carnitine Palmitoyltransferase 1C (CPT1C) in the Hippocampus. Fadó R et al. J. Biol. Chem. (2015) PubMed:26338711
AMPAR interacting protein CPT1C enhances surface expression of GluA1-containing receptors. Gratacós-Batlle E et al. Front Cell Neurosci (2014) PubMed:25698923
CPT1c is localized in endoplasmic reticulum of neurons and has carnitine palmitoyltransferase activity. Sierra AY et al. J. Biol. Chem. (2008) PubMed:18192268
Localization and effect of ectopic expression of CPT1c in CNS feeding centers. Dai Y et al. Biochem. Biophys. Res. Commun. (2007) PubMed:17559810

Access the online factsheet including applicable protocols at <https://sysy.com/product/415003> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled and biotin-labeled antibodies and control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites and hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.