

Neuropeptide S

Cat.No. 434 005; Polyclonal Guinea pig antibody, 50 µg specific antibody (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Albumin and azide were added for stabilization. For reconstitution add 50 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: not tested yet IP: not tested yet ICC: not tested yet IHC: 1 : 500 IHC-P (FFPE): 1 : 1000 up to 1 : 2000
Immunogen	Synthetic peptide corresponding to AA 70 to 89 from mouse Neuropeptide S (UniProt Id: P0C0P8)
Reactivity	Reacts with: rat (P0C0P7). Other species not tested yet.
Specificity	Recognizes Neuropeptide S. It may crossreact with the unprocessed precursor protein.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Neuropeptide S (NPS) is a 20 amino acid neuropeptide that is processed from a larger precursor. NPS was named after a conserved serine residue at its N-terminus and is found in all tetrapod vertebrates but is absent from fish.

It is predominantly expressed in the central nervous system. The distribution pattern is only partially conserved between mice and rats. In both rat and mouse brain, expression of NPS precursor mRNA is found in the locus coeruleus (LC) area. In the parabrachial area, NPS precursor mRNA is expressed in the lateral parabrachial nucleus (LPB) in the rat brain, whereas expression in the mouse brain is found in the Kölliker-Fuse (KF) nucleus. Besides these two regions, NPS precursor mRNA is also found in additional regions in the rat brain, including the principal sensory trigeminal nucleus (Pr5). NPS-responsive neurons extend projections into many other brain areas. NPS binds to its receptor NPSR1 to increase intracellular calcium concentrations. Physiologically, NPS has been characterized as a modulator of sleep/wake functions, locomotion, arousal/anxiety responses, food intake, and learning and memory.

Selected General References

Brain neuropeptide S: via GPCR activation to a powerful neuromodulator of socio-emotional behaviors. Grund T et al. Cell Tissue Res. (2019) PubMed:30112573

Anatomical characterization of the neuropeptide S system in the mouse brain by in situ hybridization and immunohistochemistry. Clark SD et al. J. Comp. Neurol. (2011) PubMed:21452235

Phylogenetic appearance of neuropeptide S precursor proteins in tetrapods. Reinscheid RK et al. Peptides (2007) PubMed:17293003

Access the online factsheet including applicable protocols at <https://sysy.com/product/434005> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies.** Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free (“no-frost”) freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at -20°C to -80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C

Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C. Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C. Adding a carrier protein (e.g., BSA) enhances long-term stability. Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C. Adding a carrier protein such as BSA improves long-term stability. Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at -20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.