

## Claudin5

Cat.No. 482 005; Polyclonal Guinea pig antibody, 50 µg specific antibody (lyophilized)

### Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Albumin and azide were added for stabilization. For <b>reconstitution</b> add 50 µl H <sub>2</sub> O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	<b>WB:</b> 1 : 1000 up to 1 : 2000 (AP staining) <b>ICC:</b> 1 : 500 (see remarks) <b>IHC:</b> not recommended <b>IHC-P:</b> 1 : 200 <b>IHC-Fr:</b> 1 : 500 up to 1 : 1000 (see remarks) <b>IHC-G:</b> 1 : 500 up to 1 : 1000 (see remarks)
Immunogen	Recombinant protein corresponding to residues near the carboxy terminus of mouse Claudin5. (UniProt Id: O54942)
Reactivity	Reacts with: rat (Q9JKD6), mouse (O54942), dog. Other species not tested yet.
Remarks	<b>ICC:</b> Methanol fixation is recommended. <b>IHC-Fr:</b> The following fixatives are possible: methanol, 4% formaldehyde/PFA. <b>IHC-G:</b> 9% glyoxal fixation is recommended.

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

### Background

Claudin5 is predominantly expressed in endothelial cells, especially brain endothelial cells (1) and therefore it is thought that the paracellular permeability of the blood–brain barrier is largely determined by the expression levels of Claudin5. An initial study using Claudin5 knockout (KO) mice clearly showed that expression of Claudin5 in the blood–brain barrier is essential for preventing the entrance of molecules with molecular weights between 400 and 800 Da in the brain. (2) The size-selective modulation of blood–brain barrier permeability has an advantage in comparison to clinically performed methods for increasing blood–brain barrier permeability, namely intra-carotid hyperosmolar mannitol administration. Mannitol administration completely disrupts the blood–brain barrier by withdrawing water from endothelial cells and enabling the entrance of bloodborne proteins into the brain. (3,4)

### Selected General References

- A molecular atlas of cell types and zonation in the brain vasculature.  
Vanlandewijck M et al. Nature (2018) PubMed:29443965
- Seizure-promoting effect of blood-brain barrier disruption.  
Marchi N et al. Epilepsia (2007) PubMed:17319915
- Size-selective loosening of the blood-brain barrier in claudin-5-deficient mice.  
Nitta T et al. J Cell Biol (2003) PubMed:12743111
- Osmotic opening of the blood-brain barrier: principles, mechanism, and therapeutic applications.  
Rapoport SI et al. Cell Mol Neurobiol (2000) PubMed:10696511

Access the online factsheet including applicable protocols at <https://sysy.com/product/482005> or scan the QR-code.



# FAQ - How should I store my antibody?

## Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

## Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies.** Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

## Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free (“no-frost”) freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at -20°C to -80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

## Product Specific Hints for Storage

### Control proteins / peptides

- Store at -20°C to -80°C

### Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C. Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C. Adding a carrier protein (e.g., BSA) enhances long-term stability. Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

### Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C. Adding a carrier protein such as BSA improves long-term stability. Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

### Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

# Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at -20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.