

Neurogenin2

Cat.No. 530 205; Polyclonal Guinea pig antibody, 50 µg specific antibody (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Albumin and azide were added for stabilization. For reconstitution add 50 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: not tested yet IP: not tested yet ICC: not tested yet IHC: 1 : 500 IHC-P: not recommended
Immunogen	Recombinant protein comprising residues near the amino and carboxy terminus of mouse Neurogenin2 (UniProt Id: P70447)
Reactivity	Reacts with: mouse (P70447). Other species not tested yet.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Neurogenins (Ngns) are basic helix-loop-helix (bHLH) transcription factors that play a crucial role in the development of the nervous system and the endocrine pancreas (1, 2). In mammals, the family comprises three members: Neurogenin1 (Ngn1), **Neurogenin2 (Ngn2)**, and Neurogenin3 (Ngn3). Ngn1 and **2** are involved in early stages of neuronal differentiation and promote the transition of neural progenitor cells into neurons and inhibit glial cell formation in a spatially and temporally regulated manner (1, 3).

Additionally, **Ngn2** plays a pivotal role in the specification of motor neurons and interneurons in the spinal cord. Phosphorylation of **Ngn2** influences its ability to direct the fate of neural progenitors, affecting the balance between different neuronal subtypes (4, 5).

Ngn3 is a pro-endocrine transcription factor, primarily associated with the development of the endocrine pancreas. It is essential for the differentiation of endocrine progenitor cells into various hormone-producing cell types within the islets of Langerhans (2).

Selected General References

bHLH transcription factors in neural development, disease, and reprogramming.
Dennis DJ et al. Brain Res (2019) PubMed:29544733

Neurogenin3: a master regulator of pancreatic islet differentiation and regeneration.
Rukstalis JM et al. Islets (2009) PubMed:21099270

Neurogenesis or neuronal specification: phosphorylation strikes again!
Lai HC et al. Neuron (2008) PubMed:18400155

Regulation of motor neuron specification by phosphorylation of neurogenin 2.
Ma YC et al. Neuron (2008) PubMed:18400164

Neurogenin1 and neurogenin2 control two distinct waves of neurogenesis in developing dorsal root ganglia.
Ma Q et al. Genes Dev (1999) PubMed:10398684

Access the online factsheet including applicable protocols at <https://sysy.com/product/530205> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies.** Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free (“no-frost”) freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at -20°C to -80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C

Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C. Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C. Adding a carrier protein (e.g., BSA) enhances long-term stability. Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C. Adding a carrier protein such as BSA improves long-term stability. Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at -20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.