

## Fibronectin

Cat.No. 550 005; Polyclonal Guinea pig antibody, 50 µg specific antibody (lyophilized)

### Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Albumin and azide were added for stabilization. For <b>reconstitution</b> add 50 µl H <sub>2</sub> O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	<b>WB:</b> 1 : 1000 <b>IP:</b> not tested yet <b>ICC:</b> 1 : 1000 <b>IHC:</b> 1 : 2000 up to 1 : 5000 (see remarks) <b>IHC-P (FFPE):</b> not tested yet
Immunogen	Full-length nativ human Fibronectin protein (UniProt Id: P02751)
Reactivity	Reacts with: human (P02751), mouse (P11276). Other species not tested yet.
Remarks	<b>IHC:</b> Antigen retrieval with citrate buffer pH 6 can be applied to improve the signal-to-noise ratio.

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

## Background

Fibronectin (FN, FN1) is an extracellular matrix glycoprotein forming a dimer (~250 kD). Two isoforms exist: plasma fibronectin from hepatocytes and cellular fibronectin made by fibroblasts, endothelial and mesangial cells (3). The FN1 gene belongs to the ECM protein family (3). Fibronectin is ubiquitous, especially in fibroblast and endothelial rich tissues, hepatocytes and renal glomerular mesangial cells (1). During embryogenesis it lines basement membranes of developing vessels and kidneys and is essential for angiogenesis; FN null mice die embryonically with severe vascular defects (2,4). In adults it participates in wound healing. By binding integrins (e.g., α5β1) and growth factors, fibronectin provides a scaffold for cell adhesion, migration and proliferation, regulating morphogenesis and angiogenesis. Clinically, elevated cellular fibronectin occurs in diabetic nephropathy and mesangial expansion (1). After cerebral ischemia it rises sharply, and an imbalance with MMP mediated degradation predicts hemorrhagic transformation (3). Altered fibronectin matrices also promote fibrosis.

## Selected General References

- Circulating fibronectin contributes to mesangial expansion in a murine model of type 1 diabetes. Klemis V et al. *Kidney Int* (2017) PubMed:28159318
- The extracellular matrix protein fibronectin promotes metanephric kidney development. Skoczynski K et al. *Pflugers Arch* (2024) PubMed:38563997
- Roles of blood-brain barrier integrins and extracellular matrix in stroke. Edwards DN et al. *Am J Physiol Cell Physiol* (2019) PubMed:30462535
- Presenilin-1 regulates the constitutive turnover of the fibronectin matrix in endothelial cells. De Gasperi R et al. *BMC Biochem* (2012) PubMed:23259730

Access the online factsheet including applicable protocols at <https://sysy.com/product/550005> or scan the QR-code.



# FAQ - How should I store my antibody?

## Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

## Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies.** Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

## Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free (“no-frost”) freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at -20°C to -80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

## Product Specific Hints for Storage

### Control proteins / peptides

- Store at -20°C to -80°C

### Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C. Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C. Adding a carrier protein (e.g., BSA) enhances long-term stability. Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

### Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C. Adding a carrier protein such as BSA improves long-term stability. Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

### Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

# Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at -20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.